



Gärtner's Epsom salts

- act to prevent signs of magnesium deficiency (needles turning brown) in pines, spruces and other conifers
- contain magnesium, which is essential for chlorophyll
- highly soluble, which means immediate availability to plants
- can also be used to supply magnesium to deciduous trees, shrubs or other woody plants
- best used in spring and late autumn

Required quantities:

- 50 –100 g/m², depending on the size of the plants
- The required quantity stated can also be mixed with the planting soil in order to enable the new plantings to have a good start
- In the event of the magnesium deficiency already being visible, as evidenced by the needles turning brown or the leaves becoming lighter in colour, improvement can be achieved by applying two doses 4 weeks apart

Application period:

Jan	Feb	March	April	May	June	July	August	Sept	Oct	Nov	Dec
-----	-----	-------	-------	-----	------	------	--------	------	-----	-----	-----

Tips:

- The effect of Epsom salts can be accelerated by dissolving them in irrigation water. The concentration in the solution should not exceed 5%. This corresponds to 50 g/l of water.
- When the Epsom salts are dissolved in the irrigation water, spreading should only take place when the weather is overcast or in the evening to avoid damage.

Dosage of Gärtner's Epsom salts:

- 1 handful approx. 40 g
- 1 tablespoon approx. 17 g

You can find the latest information on fertilisers and gardens on our website.
www.greenpartners.de

